

## Bibliografia

1. Sappey MPC. Anatomie, Physiologie, Pathologie, des VaisseauxLymphatiques Consideres chez l'Homme et les Vertebras. Paris, France: DeLaHaye Publishing; 1874.
2. Cabanas RM. An approach for the treatment of penile carcinoma. *Cancer*. 1977; 39: 456-466.
3. Morton DL, Wen DR, Wong JH, et al. Technical details of intraoperativelymphatic mapping for early stage melanoma. *Arch Surg*. 1992; 127: 392-399.
4. Giuliano AE, Kirgan DM, Guenther JM, et al. Lymphatic mapping and sentinel lymphadenectomy for breast cancer. *Ann Surg*. 1994; 220:391-398.
5. Barlin JN, Khouri-Collado F, Kim CH, et al. The importance of applying a sentinel lymph node mapping algorithm in endometrial cancer staging: beyond removal of blue nodes. *Gynecol Oncol* 2012; 125 (3): 531-5.
6. Cormier B, Diaz JP, Shih K, et al. Establishing a sentinel lymph node mapping algorithm for the treatment of early cervical cancer. *Gynecol Oncol*. 2011 Aug;122:275-280.
7. Lecuru F, Mathevet P, Querleu D, et al. Bilateral negative sentinel nodes accurately predict absence of lymph node metastasis in early cervical cancer: results of the SENTICOL study. *J Clin Oncol* 2011; 29:1686-91.
8. Koh WJ, Greer BE, Abu-Rustum NR, et al. Cervical Cancer Version 2.2015. *J Natl Compr Canc Netw* 2015; 13 (4): 395-404.
9. Ercoli A, Dalmas V, Iannone V, et al. The lymphatic drainage of the uterine cervix in adult fresh cadavers: anatomy and surgical implications. *Eur J Surg Oncol* 2010; 36 (3): 298-303.
10. Horn LC, Hentschel B, Fischer U, et al. Detection of micrometastases in pelvic lymph nodes in patients with carcinoma of the cervix uteri using step sectioning: Frequency, topographic distribution and prognostic impact.*Gynecol Oncol*. 2008; 111 (2): 276-81.
11. Lécuru F, Mathevet P, Querleu D, et al. Bilateral negative sentinel nodes accurately predict absence of lymph node metastasis in early cervical cancer: results of the SENTICOL study. *J Clin Oncol*. 2011; 29(13): 1686-91.
12. Bats AS, Mathevet P, Buenerd A, et al. The sentinel node technique detects unexpected drainage pathways and allows nodal ultrastaging in early cervical cancer: insight from the multicenter prospective SENTICOL study). *Ann Surg Oncol* 2013; 20: 413-22.
13. Wang XJ, Fang F, Li YF. Sentinel-lymph-node procedures in early stage cervical cancer: a systematic review and meta-analysis. *Med Oncol*. 2015; 32 (1): 385.
14. Altgassen C, Hertel H, Brandstädt A, et al. Multicenter validation study of the sentinel lymph node concept in cervical cancer: AGO Study Group. AGO Study Group. *J Clin Oncol*. 2008; 26(18): 2943-51.
15. Eriksson LR, Covens A. Sentinel lymph node mapping in cervical cancer: the future? *BJOG*. 2012 Jan;119(2):129-33.
16. Hoogendam JP, Veldhuis WB, Hobbelink MG, et al. 99mTc SPECT/CT Versus Planar Lymphoscintigraphy for Preoperative Sentinel Lymph NodeDetection in Cervical Cancer: A Systematic Review and Metaanalysis. *J Nucl Med*. 2015; 56 (5): 675-80.
17. Buda A, Elisei F, Arosio M, et al. Integration of hybrid single-photon emission computed tomography/computed tomography in the preoperative assessment of sentinel node in patients

- with cervical and endometrial cancer: our experience and literature review. *Int J Gynecol Cancer*. 2012; 22 (5): 830-5.
18. Jewell EL, Huang JJ, Abu-Rustum NR, et al. Detection of sentinel lymph nodes in minimally invasive surgery using indocyanine green and near-infrared fluorescence imaging for uterine and cervical malignancies. *Gynecol Oncol*. 2014; 133 (2): 274-7.
  19. Cibula D, Abu-Rustum NR, Dusek L, et al. Prognostic significance of low volume sentinel lymph node disease in early-stage cervical cancer. *Gynecol Oncol*. 2012; 124 (3): 496-501.
  20. Cibula D, Zikan M, Slama J, et al. Risk of micrometastases in non-sentinel pelvic lymph nodes in cervical cancer. *Gynecol Oncol*. 2016; 143 (1): 83-6.
  21. Buda A, Di Martino G, Vecchione F, et al. Optimizing Strategies for Sentinel Lymph Node Mapping in Early-Stage Cervical and Endometrial Cancer: Comparison of Real-Time Fluorescence With Indocyanine Green and Methylene Blue. *Int J Gynecol Cancer*. 2015; 25 (8): 1513-8.
  22. Buda A, Bussi B, Di Martino G, et al. Sentinel Lymph Node Mapping With Near-Infrared Fluorescent Imaging Using Indocyanine Green: A New Tool for Laparoscopic Platform in Patients With Endometrial and Cervical Cancer. *J Minim Invasive Gynecol*. 2016 Feb 1;23(2):265-9.
  23. Buda A, Dell'Anna T, Vecchione F, et al. Near-Infrared Sentinel Lymph Node Mapping With Indocyanine Green Using the VITOM II ICG Exoscope for Open Surgery for Gynecologic Malignancies. *J Minim Invasive Gynecol*. 2016 May-Jun;23(4):628-32.
  24. Buda A, Crivellaro C, Elisei F, et al. Impact of Indocyanine Green (ICG) for sentinel node mapping in early stage endometrial and cervical cancer: comparison with conventional radiotracer TC99m and/or blue dye. *Ann Surg Oncol* 2016; 23 (7): 2183-91.
  25. Buda A, Elisei F, Palazzi S, et al. Quality of care for cervical and endometrial cancer patients: the impact of different techniques of sentinel lymph node mapping on Patient Satisfaction. *Ann Surg Oncol* 2016; 23 (9): 2975-81.
  26. Buda A, Papadia A, Zapardiel I, et al. From conventional radiotracer Tc-99(m) with blue dye to indocyanine green fluorescence: a comparison of methods towards optimization of sentinel lymph node mapping in early stage cervical cancer for a laparoscopic approach. *Ann Surg Oncol*. 2016; 23 (9): 2959-65.